

Continuum effects in reactions involving weakly bound nuclei

A. M. Moro^a, R. Crespo^a, F. Nunes^b and I. J. Thompson^c

^aDepartamento de Física, Instituto Superior Técnico, 1049-001 Lisboa, Portugal

^bUniversidade Fernando Pessoa, Praça 9 de Abril, 4200 Porto, Portugal

^cDepartment of Physics, University of Surrey, Guildford GU2 5XH, UK

In this contribution we investigate the role of the continuum spectrum in reactions of astrophysical interest. In particular, the influence of these coupling effects in the description of elastic and transfer reactions is discussed. We examine the validity of the Distorted Wave Born Approximation (DWBA) as a tool to extract the astrophysical $S_{17}(0)$ factor from the transfer reaction $^{14}\text{N}(^7\text{Be}, ^8\text{B})^{13}\text{C}$. For this purpose, we present calculations for this reaction comparing the DWBA method with the CDCC-Born approximation, in which continuum effects are explicitly incorporated.

1. INTRODUCTION

In recent years, much of the effort in nuclear physics has been devoted to the study of nuclear reactions of astrophysical interest. The direct measurement of these reactions at the relevant energies (tens of keV or less) is still beyond the capability of the existing facilities. From the theoretical point of view, the description of these reactions can be substantially simplified by noting that, at these low energies, the reaction mechanism is only sensitive to the tail of the wave-function for the participants. Thus, the rate of the capture reaction $a(p, \gamma)b$ is essentially determined by the asymptotics of the single-particle overlap $(b|a)$ which, in turn, is proportional to a Whittaker function. The proportionality constant, known as the asymptotic normalization coefficient (ANC), determines unambiguously the capture cross section at zero energy.

This feature constitutes the basis of the ANC method, which extracts the ANC for the overlap $(b|a)$ from the analysis of a transfer reaction $A(a, b)B$. As an important prerequisite, the selected transfer reaction has to be peripheral. Under this condition, the DWBA amplitude for the transfer process is proportional to the product of the ANC's C_{ba} and C_{BA} . Thus, by comparison of the calculated and measured transfer cross section, it is possible to extract the value for C_{ba} , provided that the ANC C_{BA} is already known. Due to the approximations involved in the DWBA, the applicability of the method requires that the transfer occurs in one-step and that the entrance and exit channels are well described by phenomenological optical potentials [1].

When weakly bound nuclei are involved, the transfer mechanism might occur through continuum states. Thus, it is not obvious that these multi-step processes are properly accounted for by an optical model potential, as it is assumed in the DWBA approach. In




Figure 1. Angular distribution for the differential elastic cross sections calculated with the CDCC method (thick solid line) and RDF potential (thin solid line). The dashed line is the CDCC calculation without continuum couplings.

these situations, it may result more adequate to treat explicitly the couplings involving the continuum states. A suitable formalism to include these effects is the CDCC-BA approach. Here, the partition that contains the weakly bound nucleus is described within a model space that includes the continuum spectrum of that nucleus, by means of a procedure of discretization. Formally, this is done by replacing the total wavefunction in the exact transition amplitude by a solution of a Continuum Discretized Coupled Channels calculation [2].

Recently, the reaction $^{14}\text{N}(^7\text{Be},^8\text{B})^{13}\text{C}$ at a bombarding energy of 85 MeV has been measured at the Texas A&M Cyclotron facility in order provide an indirect determination of the astrophysical S_{17} factor, using the ANC for the overlap $(^7\text{Be},^8\text{B})$. This ANC was extracted by analyzing the experimental data within the DWBA approach [3]. Since the ^8B nucleus is weakly bound ($\epsilon_0 = 0.137$ MeV), transfer could proceed through the ^8B continuum states and hence this approach could be inappropriate. In this work, we aim to evaluate the effect of the ^8B continuum in the cited transfer reaction. For this purpose, we have performed an analysis of the data using the CDCC-BA approach and comparing our results with those obtained in the DWBA method. Since the DWBA and CDCC-BA approaches differ essentially in the description of the exit channel, we start by analyzing the elastic scattering $^8\text{B}+^{13}\text{C}$.

2. ANALYSIS OF THE ENTRANCE AND EXIT CHANNELS

In the DWBA analysis of Azhari et al [3], the elastic scattering $^7\text{Be}+^{14}\text{N}$ was described using a double folding potential, with an average normalization constant extracted from the analysis of reaction with nearby nuclei [4]. As shown in [3], this renormalized double

folding (RDF) potential reproduces very well the experimental data for the elastic scattering ${}^7\text{Be}+{}^{14}\text{N}$. In the present analysis, we have also adopted this potential to describe the entrance channel in both the DWBA and CDCC-BA calculations.

The lack of experimental data for the reaction ${}^8\text{B}+{}^{13}\text{C}$, due to the small intensities of the present beams of ${}^8\text{B}$, makes the optical potential for the exit channel more uncertain. In [3], the RDF potential was used. Note that this ambiguity is absent in the CDCC approach, since the ${}^8\text{B}+{}^{13}\text{C}$ interaction is described in terms of the cluster-target interactions. In the present work, we have assumed that the ${}^8\text{B}$ nucleus consists of an inert ${}^7\text{Be}$ core plus a valence proton. The ${}^7\text{Be}-p$ continuum is discretized into $N = 10$ bins, uniformly spaced in energy, and up to $\epsilon_{\text{max}} = 9$ MeV. We took into account s , p , and d partial waves for the relative motion. Details of binding and optical potentials used in the calculations can be found in [2] and references therein.

In Fig. 1 we show the calculated elastic scattering for ${}^8\text{B}+{}^{13}\text{C}$ at $E_{\text{lab}}=78$ MeV. The thin solid line is the calculation with the RDF potential, whereas the thick solid line is the full CDCC result. We find a good agreement between both calculations at small angles, but they differ significantly at larger angles, in the region where continuum effects become relevant. Although experimental data would be required to clarify the accuracy of both methods, this result casts doubt on the validity of global parameterizations when weakly bound nuclei are involved. Also shown in this figure (dashed line) is the CDCC calculation without continuum couplings, which corresponds just to a cluster folding calculation. By comparing this calculation with the full CDCC, we find that continuum effects become important for scattering angles above 30 degrees. Although not shown in this figure, we found also that the calculation with only g.s.-continuum couplings gives a result very close to the full CDCC, indicating that continuum-continuum effects are small in this reaction.

3. THE TRANSFER REACTION

Although the DWBA and CDCC-BA amplitudes differ in the description of the exit channel, they involve the same transition operator. In prior form, this operator is $V_{p,14\text{N}} + V_{\text{rem}}$, where $V_{\text{rem}} = U_{7\text{Be},13\text{C}} - U_{7\text{Be},14\text{N}}$ is commonly referred as remnant term.

In order to allow a reliable comparison between the DWBA and CDCC-BA calculations, the same core-core (${}^7\text{Be} + {}^{13}\text{C}$) interaction was used in both methods. In particular, we took the ${}^7\text{Li}+{}^{13}\text{C}$ optical potential derived in [4]. The $({}^{14}\text{N}|{}^{13}\text{C})$ overlap was described using a $p_{1/2}$ single-particle wave-function, with spectroscopic factor 0.604 [5]. For the $({}^8\text{B}|{}^7\text{Be})$ overlap, we assumed a pure $p_{3/2}$ configuration with spectroscopic factor $S_{p_{3/2}} = 0.733$. The CDCC calculations had to be multiplied also by the geometric factor $5/16$, that arises from the fact that our model assumes that ${}^7\text{Be}$ has zero spin [2].

In Fig. 2 we present the DWBA transfer cross sections as a function of the total angular momentum, L , and the impact parameter $R = L/k_0$, where k_0 is the incident linear momentum. It is observed that the transfer occurs mainly at distances $R > R_1 + R_2$, where R_1 and R_2 are an estimate for the radii of the projectile and target. This confirms the peripheral character of the present reaction, which is a prerequisite for the applicability of the ANC method.

In Fig. 3 the experimental and calculated cross sections for the transfer reaction are shown. The thin solid line and the thick solid line correspond to the DWBA and CDCC-

BA calculations, respectively. The dotted line is a CDCC-BA calculation without continuum couplings. We see that the three curves give very similar results, indicating that in this reaction continuum effects are not important and thus supporting the validity of the DWBA as a reliable method to extract the ANC for the overlap (${}^8\text{B}|{}^7\text{Be}$). Also shown in this figure, is the DWBA calculation without remnant term in the transition amplitude. The sizable difference with the exact DWBA calculation shows that the full DWBA transition operator is essential for a quantitative analysis of this reaction.




Figure 2. Transfer cross section versus the total angular momentum. The shaded region corresponds to $R < R_1 + R_2$ where R_1 and R_2 are estimates for the radii of the colliding nuclei.




Figure 3. Experimental and calculated transfer cross sections for the reaction ${}^{14}\text{N}({}^7\text{Be}, {}^8\text{B}){}^{13}\text{C}$ at 84 MeV.

In order to understand the reason why continuum effects have such a small effect on the transfer cross sections we compare in Fig. 4 the DWBA cross section for the process ${}^{14}\text{N}({}^7\text{Be}, {}^8\text{B}){}^{13}\text{C}$ with the breakup cross section (calculated with CDCC) for the exit channel ${}^8\text{B}+{}^{13}\text{C}$. In order to have enhanced continuum effects in ${}^{14}\text{N}({}^7\text{Be}, {}^8\text{B}){}^{13}\text{C}$, both the continuum excitation couplings and the transfer couplings need to be strong. In Fig. 4(a) we see that, as expected, the first condition is satisfied in this reaction, since there is an important breakup probability to continuum states. However, transfer couplings to continuum states are comparatively much smaller (partially due to the unfavourable Q value), as becomes evident in Fig. 4(b). Thus, the multi-step process consisting on transfer to the ${}^7\text{Be}+p$ continuum and deexcitation to the ${}^8\text{B}$ ground state are suppressed.

4. CONCLUSIONS

In summary, we have studied the effect of ${}^8\text{B}$ two-body continuum in the transfer reaction ${}^{14}\text{N}({}^7\text{Be}, {}^8\text{B}){}^{13}\text{C}$. For this purpose, we have compared the standard DWBA method,




Figure 4. (a) Breakup cross section, calculated in the CDCC approach, for the scattering ${}^8\text{B}+{}^{13}\text{C}$ at 74 MeV. (b) DWBA cross section leading to ${}^8\text{B}$ continuum states for the reaction ${}^{14}\text{N}({}^7\text{Be}, {}^8\text{B}^*){}^{13}\text{C}$ at 84 MeV.

with the CDCC-BA approach. The later takes into account ground state–continuum couplings and continuum–continuum couplings for the ${}^8\text{B}$ continuum.

We find that the elastic scattering predicted by the CDCC calculation for the exit channel (${}^8\text{B}+{}^{13}\text{C}$) agrees very well with the renormalized double folding potential used in [4] at small angles, but differs significantly at larger angles. However, these discrepancies do not seem to have any effect in the forward angle transfer cross sections. Therefore, our calculations confirm the applicability of the DWBA for the present reaction and, hence, the value of the ANC extracted in [3].

ACKNOWLEDGMENTS

This work has been partially supported by Fundação para a Ciência e a Tecnologia (F.C.T.) under Grant No. SAPIENS/36282/99 and EPSRC under Grant No. GR/M/82141. One of the authors, A.M.M., acknowledges a F.C.T. post-doctoral grant.

REFERENCES

1. J. C. Fernandes, R. Crespo, and F. M. Nunes, Phys. Rev. **C61** (2000) 064616.
2. A. M. Moro, R. Crespo, F. Nunes, and I. J. Thompson, Phys. Rev. **C66** (2002) 024612.
3. A. Azhari *et al.*, Phys. Rev. **C60** (1999) 055803.
4. L. Trache *et al.*, Phys. Rev. **C61** (2000) 024612.
5. L. Trache *et al.*, Phys. Rev. **C58** (1998) 2715.